INTRATUMORAL INJECTION OF LTX-315 IN COMBINATION WITH PEMBROLIZUMAB IN PATIENTS WITH ADVANCED MELANOMA REFRACTORY TO PRIOR PD-1/PD-L1 THERAPY: INTERIM RESULTS FROM THE ATLAS-IT-05 TRIAL

STÉPHANE DALLE¹, MIGUEL F SANMAMED², JOHN M KIRKWOOD³, CAROLINE ROBERT⁴, MARTA NYAKAS⁵, THOMAS MARRON⁶, LAURENT MORTIER⁷, GRAEME CURRIE⁸, VIBEKE SUNDVOLD⁸, ØYSTEIN REKDAL⁸, BALDUR SVEINBJØRNSSON⁸, JACKIE EARABINO⁸, KARSTEN BRUINS SLOT⁸, ADI DIAB⁹

SPAIN; 3 UNIVERSITY OF PITTSBURGH MEDICAL CENTRE - HILMAN CANCER CENTRE, USA; 4 GUSTAVE ROUSSY/PARIS SACLAY UNIVERSITY, FRANCE;

¹ CHU HÔPITAL LYON SUD, FRANCE; ² CLINICA UNIVERSIDAD DE NAVARRA, ⁵ OSLO UNIVERSITY HOSPITAL-RADIUMHOSPITAL, NORWAY; ⁶ ICAHN SCHOOL OF MEDICINE AT MOUNT SINAI, USA;7 CHU LILLE, FRANCE; 8 LYTIX BIOPHARMA AS, NORWAY; 9 MD ANDERSON CANCER CENTRE, USA



1051P

BACKGROUND

LTX-315 is a first-in-class oncolytic peptide of non-viral origin that is in development for intratumoral treatment of solid tumors (1,2)


Pre-clinical studies of LTX-315 demonstrate:

- Unique immunogenic cell death mode of action by causing mitochondrial lysis and disintegration of cytoplasmic organelles resulting in effective release of danger signals and a broad repertoire of tumor antigens (3-6)
- Reduced number of immunosuppressive cells (T reg and myeloid derived suppressor cells) (7)
- Enhanced infiltration of T cells and T cell clonality (8)
- Complete regression of injected and non-injected tumors (i.e. systemic immune response) (8-10)

Clinical studies of LTX-315 demonstrate:

- Enhanced infiltration of T cells and T cell clonality (11, 12)
- Regression of injected and non-injected tumors (i.e. systemic immune response) (12)
- Generation of tumor-specific T cells (13)

LTX-315 - UNIQUE MODE OF ACTION RESULTS IN EFFECTIVE RELEASE OF POTENT IMMUNOSTIMULANTS AND ANTIGENS



STUDY OBJECTIVES AND ENDPOINTS

Objectives

Evaluate the efficacy and safety of intratumoral LTX-315 in combination with pembrolizumab in patients with Stage IIIB-IVm1b melanoma, who have progressed on or after prior treatment with a PD-1/PD-L1 inhibitor

Primary Efficacy Endpoint


- Objective Response Rate (ORR) using RECIST v1.1 criteria assessed by investigators
- Disease Control Rate (DCR) using RECIST v1.1 criteria assessed by investigators

Secondary Efficacy Endpoint

- Regression of injected lesions assessed by CT/MRI or ultrasound measurements by
- Incidence and severity of treatment emergent adverse events related to LTX-315

STUDY DESIGN

Open label, single-arm, phase II study with 10 study sites in Europe and the US (NCT04796194)



KEY INCLUSION AND EXCLUSION CRITERIA

- Histologically confirmed, Stage IIIB-IVm1b unresectable melanoma
- Confirmed disease progression on or after prior treatment with PD-1/PD-L1 inhibitor
- ≤3 prior lines of systemic treatment for metastatic disease
- ECOG performance status of 0-1
- At least 1 superficial, non-visceral tumor lesion accessible for injection superficial lymph nodes with metastatic disease can also be injected
- LDH ≤ 2 x ULN
- No ocular or mucosal melanoma diagnosis

PATIENT DISPOSITION

	Number of patients
Patients with available data at cutoff date (13 September 2023)	20
Patients with melanoma diagnosis	19
Patients included in Safety Analysis Set (SAS)	20
Patients included in Efficacy Analysis Set (EAS)	14
 Reasons for exclusion from EAS No available post-baseline scan at cutoff date yet Patient withdrew consent prior to first post-baseline scan Only 1 available post-baseline scan and insufficient follow-up No melanoma diagnosis* 	3 1 1 1

Patient with acinic cell carcinoma enrolled under earlier protocol version

Median duration on study was 15 weeks at cutoff date

BASELINE CHARACTERISTICS

Baseline characteristic	All patients (N=20)	Baseline characteristic	All patients (N=20)
Mean age (range)	68 years (42-91)	Prior systemic treatment lines in metastatic setting	
Sex Female Male	9 (45%) 11 (55%)	0 1 2 3	1 (5%)** 11 (55%) 6 (30%) 2 (10%)
ECOG 0 1	16 (80%) 4 (20%)	Prior lines of treatment with checkpoint inhibitor	1 (5%)**
Melanoma stage Stage IIIB	1 (5%)	1 ≥2	6 (30%) 13 (65%)
Stage IIIC Stage IIID Stage IVm1a Stage IVm1b	4 (20%) 2 (10%) 7 (35%) 5 (25%)	BRAF status Positive Negative	5 (25%) 15 (75%)
Prior systemic treatment for metastatic disease*		Negative	13 (7370)
BRAF/MEK PD-(L)1 monotherapy PD-(L)1+ CTLA-4 PD-(L)1 + other CTLA-4+ other	4 (20%) 10 (50%) 8 (40%) 6 (30%) 1 (5%)	LDH Normal >ULN	9 (45%) 11 (55%)


*More than 100% as some patients received >1 systemic metastatic treatment line All melanoma patients had prior surgery

BEST OVERALL RESPONSE - RECIST V1.1


Best overall response (RECIST v1.1)	N (%)
Complete response	0 (0%)
Partial Response*	1 (7%)
Stable Disease	5 (36%)
Progressive Disease	8 (57%)

Objective Response Rate (ORR) = 7% (95% CI 1-30%) Disease Control Rate (DCR)= 43% (95% CI 20-70%)


BEST CHANGE IN RECIST TARGET LESIONS



CHANGE IN RECIST TARGET LESIONS



RESPONSE ASSESSMENTS PER RECIST V1.1



RESPONSE IN INJECTED LESIONS

- 9 out of 21 (43%) evaluable injected lesions showed complete regression by CT scan as best response after start of treatment*
- Complete regression was shown in 3 out of 11 (27%) evaluable patients by CT scan *Analysis includes EAS population, except: Injected lesions that were not injected per injection plan



perpendicular diameters

12 mm

Injected lesions without baseline assessment of lesion size





EVENTS (>10%) IN SAS

OVERVIEW OF TREATMENT EMERGENT ADVERSE

Preferred AE term	Patients n (%)
Injection site pain	15 (75%)
Asthenia	5 (25%)
Pruritus	5 (25%)
Anemia	5 (25%)
Fatigue	4 (20%)
Injection site erythema	4 (20%)
Injection site swelling	3 (15%)
Hypertension	3 (15%)

LTX-315 TREATMENT-RELATED ADVERSE EVENTS (>10%) IN SAS

Preferred AE term	Grade 1-2	Grade 3	Grade 4	Grade 5	Patients - n (%)
Injection site pain	10 (50%)	5 (25%)	0	0	15 (75%)
Injection site erythema	3 (15%)	1 (5%)	0	0	4 (20%)
Injection site swelling	3 (15%)	0	0	0	3 (15%)

- The most common (>10%) LTX-315 treatment-related adverse events were related to injections and mostly mild, self-limiting and manageable in clinical practice.
- There was no increase in immune-related adverse events.
- No grade 4-5 treatment-related adverse events were reported

CASE - MELANOMA PATIENT WITH CLINICALLY RELEVANT SYSTEMIC RESPONSE

- 75-year-old male with Stage IVm1a, nodular melanoma (BRAF positive)
- Multiple metastases in lymph nodes and gluteal muscle at baseline
- Prior treatment with nivolumab (adjuvant setting) and BRAF/MEK inhibitor
- Treated with in total 20 intratumoral LTX-315 injections in 4 lesions on prescribed dosing days and 2 cycles (200 mg) + 3 cycles (400 mg) pembrolizumab
- Non-injected RECIST target lesion in left gluteal muscle

NON-INJECTED LESION


(metastatic setting)

BASELINE

DAY 43

Superficial lesion located in gluteus muscle

Partial response as best overall response at cutoff date with RECIST target lesion shrinkage of 89%



measurements based on caliper/ultrasound from Day 1-29; thereafter on CT scan.

CONCLUSION

- Intratumoral treatment with LTX-315 is well-tolerated with generally mild to moderate treatment-related adverse events.
- Adverse events related to the intratumoral injections were self-limited and easily manageable in clinical practice.
- The combination regimen showed preliminary signs of tumor shrinkage and stabilization in heavily pre-treated patients with PD-1/PD-L1 inhibitor refractory metastatic melanoma - enrolled patients had generally poor prognostic factors and some patients had also failed BRAF/MEK inhibition.
- The efficacy signal is encouraging with a disease control rate of 43% and 1 patient achieving a partial response to date.
- There is evidence of tumor shrinkage in both injected and in non-injected lesions.
- The trial is currently ongoing, and data are still evolving and considered immature at this stage - further details will be shared in a future presentation.

REFERENCES

1. Haug, B.E. et al; Journal of Medicinal 9. Camilio, K. et al.; Oncoimmunology Chemistry (2016).

10. Nestvold, J. et al.; Oncoimmunology

11. Jebsen, N et al.; Journal of Medical

12. Spicer, J. et al.; Clin Cancer Res

13. Nielsen, M et al. Journal of Clinical

Case Reports (2019).

Oncology (2022).

- 2. Sveinbjørnsson, B. et al.; Future Medicinal Chemistry (2017).
- 3. Zhou, H. et al.; Oncotarget (2015).
- 4. Eike, L-M. et al.; Oncotarget (2015).
- 5. Forveille, S. et al.; Cell Cycle (2015). 6. Zhou, H. et al.; Cell Death Disease
- 7. Yamazaki, T. et al.; Cell Death and Differentiation (2016).
- 8. Camilio, K. et al.; Cancer Immunology Immunotherapy (2014).

Dol Stephane Dalle: research grants and advisory board participation fees paid to institution by MSD, BMS, Pierre Fabre; SD spouse is a Sanofi employee.

Patient accrual (SAS): CHU Lyon 5; UPMC 3; MD Anderson 3; CU Navarra 3; CHRU Lille 2; Radiumhospital 2; Gustave Roussy 1; Mount Sinai 1.

Contact info: post@lytixbiopharma.com