18P - Trastuzumab in HER2-amplified or overexpressed gallbladder cancer: Experience from an Indian cancer center



D. Mondal, S. Ganguly, J. Ghosh, P. Pandey, S. Roy, B. Biswas Tata Medical Center, Kolkata, India

INTRODUCTION

The prognosis of advanced gall bladder carcinoma (GBC) remains poor despite the addition of newer targeted therapies. HER2 is a potential therapeutic target expressed in up to 20% GBC.

We present an analysis of advanced or metastatic HER2-amplified or overexpressed GBC treated with trastuzumab and chemotherapy from an Indian cancer center.

METHODS

- Retrospective review was conducted for patients with advanced or metastatic GBC with immunohistochemical evidence of HER2 overexpression (IHC 3+) or HER2 amplification detected with FISH, treated with trastuzumab and first-line chemotherapy between 2020 and 2021.
- Outcome measures were objective response rate (ORR), progression-free survival (PFS), overall survival (OS), and adverse events attributable to trastuzumab.

- recurred after prior curative treatment.
- oxaliplatin as chemotherapy.

- were attributable to trastuzumab.

RESULTS

• Twenty-six patients (age range, 44-71) with GBC were treated with trastuzumab and chemotherapy. • Twenty patients (76.9%) had newly diagnosed advanced disease while six patients (23.1%) had

• Twenty-four patients (92.3%) received gemcitabine-cisplatin and two received (7.7%) gemcitabine-




Fig 1. Response Rate

• Partial response was seen in 11 patients (42.3%) while 10 patients (38.5%) had stable disease (objective response rate 42.3%, overall response rate 80.8%) [Fig 1].

• At a median follow-up of 15.7 months, median PFS was 9.7 months (95% CI, 9.3-10) and median OS was 11.9 months (95% CI, 11.2-12.5) [Fig 2 and Fig 3].

• Grade 2 ejection fraction drop was seen in 1 patient (3.8%). No adverse events of grade 3 or more






Fig 2. Progression-free Survival




Fig 2. Overall Survival

REFERENCES

Das CK, Patel A, Raj A, Gupta VG, Mehta P, Bhethanabhotla S, et al. Anti-Her2neu directed therapy in advanced gall bladder cancer: A prospective, multicenter experience from India. JCO 2020;38:e16682-e16682.

Jeong H, Jeong JH, Kim K-P, Lee SS, Oh DW, Park DH, et al. Feasibility of HER2-Targeted Therapy in Advanced Biliary Tract Cancer: A Prospective Pilot Study of Trastuzumab Biosimilar in Combination with Gemcitabine Plus Cisplatin. Cancers (Basel) 2021;13:161.

DISCLOSURES

No conflicts of interest to declare

Email: dr.debapriyo89@gmail.com

Twitter: @debonco